

TOTAL SOLUTIONS

QUICK STRIKE INSECTICIDE

- CLEAR-STABLE - LEAVES NO OILY RESIDUES
- WATER-BASED, FAST ACTING INSECTICIDE
- KILLS AND REPELS BITING FLIES
- FOR USE IN INSTITUTIONS, RESTAURANTS, FOOD PROCESSING PLANTS, DAIRIES, LIVESTOCK AND OUTDOORS
- FOR CONTROL OF FLEAS AND BROWN DOG TICKS ON PREMISES AND ON PETS
- KILLS CRAWLING AND FLYING INSECTS ON CONTACT
- CONTAINS NATURAL [PYRETHRIN] [INSECTICIDE] DERIVED FROM FLOWERS

ACTIVE INGREDIENTS:

Pyrethrins 0.10%
 * Piperonyl Butoxide, Technical 1.00%

INERT INGREDIENTS: 98.90%
 TOTAL 100.00%

* Equivalent to min. 0.8% (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for first aid instructions and additional precautionary statements.

EPA REG NO. 10088-96-68562 • EPA EST NO. 10088-WI-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before space spraying or surface applications.

FIRST AID

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated. **IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. For terrestrial use, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

GENERAL USE DIRECTIONS

INDOOR APPLICATION:

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or

covered during treatment. All food processing surfaces should be covered during treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

SURFACE SPRAYING: To control *Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clovermites, Cheese Mites, Earwigs, Sowbugs, Boxelder Bugs, Centipedes, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles:* Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For *Silverfish*, spray bookcases. For *Ants*, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors, windows and transoms and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of *House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indianmeal Moths and Tobacco Moths,* direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON LIVESTOCK: To protect cattle, horses and hogs from *Horn Flies, House Flies, Mosquitoes* and *Gnats* apply a light mist sufficient to wet the surface of the hair.